

## 新元古代南华盆地Sturtian冰川沉积物中碳同位素负偏及其地质意义

王萍, 王健, 杜远生, 周琦, 余文超, 徐源, 袁良军, 潘文, 徐亚军

## 引用本文:

王萍, 王健, 杜远生, 等. 新元古代南华盆地Sturtian冰川沉积物中碳同位素负偏及其地质意义[J]. 沉积学报, 2024, 42(6): 2020-2030.

WANG Ping, WANG Jian, DU YuanSheng, et al. Negative Carbon Isotope Excursions of the Statian Glacial Sediments in the Neoproterozoic Nanhua Basin and Their Geological Significance[J]. Acta Sedimentologica Sinica, 2024, 42(6): 2020-2030.

## 相似文章推荐(请使用火狐或IE浏览器查看文章)

## Similar articles recommended (Please use Firefox or IE to view the article)

## 中国华南渝东北城口地区下寒武统烃源岩发育环境与形成机制

Development Environment and Formation Mechanism of Lower Cambrian Source Rocks in the Chengkou Area, Northeast Chongqing, South China

沉积学报. 2020, 38(5): 1111-1122 https://doi.org/10.14027/j.issn.1000-0550.2019.102

## 松辽盆地晚白垩世陆表古温度定量重建

Quantitative Paleotemperature Reconstruction of Late Cretaceous Nenjiang Formation in Songliao Basin: A case study of the LD67 Core 沉积学报. 2020, 38(4): 759–770 https://doi.org/10.14027/j.issn.1000–0550.2019.079

## 小冰期气候的湖北石笋碳同位素记录

Carbon Isotope Record in Stalagmites from Hubei during the Little Ice Age 沉积学报. 2020, 38(3): 497–504 https://doi.org/10.14027/j.issn.1000–0550.2019.058

## 赣西北下寒武统观音堂组碳硅泥岩地球化学特征及其沉积环境演化

Geochemical Characteristics and Sedimentary Environment Evolution of the Carbonaceous–Siliceous Mudstone of the Guanyintang Formation of the Lower Cambrian, Northwestern Jiangxi Province

沉积学报. 2019, 37(2): 278-291 https://doi.org/10.14027/j.issn.1000-0550.2018.108

## 湖南茶陵盆地晚白垩世古降水氧同位素

Late Cretaceous Oxygen Isotope of Paleoprecipitation in Chaling Basin, Hunan Province 沉积学报. 2018, 36(6): 1169–1176 https://doi.org/10.14027/j.issn.1000–0550.2018.107

文章编号:1000-0550(2024)06-2020-11

DOI: 10.14027/j.issn.1000-0550.2023.140

# 新元古代南华盆地 Sturtian 冰川沉积物中碳同位素 负偏及其地质意义

王萍1,王健1,杜远生2,周琦3,余文超2,徐源2,袁良军4,潘文4,徐亚军2

1.河南理工大学资源环境学院,河南焦作 454003

2.中国地质大学(武汉)地球科学学院,武汉 430074

3. 自然资源部基岩区矿产资源勘查工程技术创新中心,贵阳 550081

4.贵州省地质矿产勘查开发局103地质大队,贵州铜仁 554300

摘 要【目的】新元古代Sturtian冰期(~717~660 Ma)在全球范围内广泛发育,在南华盆地中也有良好的记录,但是目前对冰期的碳循环过程研究较少。为了解Sturtian冰期期间的碳循环过程,对冰期期间沉积的铁丝坳组碳同位素进行系统的研究。【方法】 以黔东松桃地区高地ZK2115钻孔为研究对象,对铁丝坳组进行高分辨率的有机碳和无机碳同位素(δ<sup>13</sup>C<sub>eat</sub>和δ<sup>13</sup>C<sub>eat</sub>)测试分析。 【结果】铁丝坳组δ<sup>13</sup>C<sub>eat</sub>值介于-9.29‰~-3.37‰(平均值为-7.24‰),具有明显的负偏,δ<sup>13</sup>C<sub>eat</sub>值介于-33.63‰~-23.35‰(平均值为 -29.29‰)。此外,δ<sup>13</sup>C<sub>eat</sub>和δ<sup>13</sup>C<sub>eat</sub>具有明显的正相关关系,表明地层中的无机碳同位素组成不受成岩作用的影响,δ<sup>13</sup>C<sub>eat</sub>代表了 Sturtian冰期期间南华盆地中溶解无机碳(Dissolved Inorganic Carbon, DIC)库的碳同位素组成,并且地层中的有机质主要是通过 光合作用形成的,有机质中固定的碳主要来源于海水中的 DIC库。【结论】结合已发表的铁丝坳组总有机碳(Total Organic Carbon, TOC)数据(平均值为0.2%),这意味着在冰期这种极端气候条件下,光合作用虽然仍在进行,但是其速率比较缓慢,产生有机质 的量相对较少。对Sturtian冰期沉积物中碳同位素组成的研究,有助于进一步探讨冰期的碳循环过程,同时也为深入理解极端气 候条件下的古海洋环境及生物地球化学循环提供依据。

关键词 Sturtian冰期;铁丝坳组;碳同位素;南华盆地

**第一作者简介** 王萍,女,1991年出生,博士,讲师,沉积地质学,E-mail: wangping\_2016@163.com 通信作者 杜远生,男,教授,E-mail: duyuansheng126@126.com

中图分类号 P534.3 文献标志码 A

0 引言

新元古代记录了两次全球性的冰期事件,即 Sturtian冰期(~717~660 Ma)和Marinoan冰期(~654~ 635 Ma)<sup>[1:3]</sup>。"雪球地球"假说<sup>[46]</sup>认为在这两次冰期事 件过程中,地球表面完全被冰川覆盖,生态系统崩 塌,水循环停滞,并且大陆风化作用基本停止<sup>[5:6]</sup>。随 着对雪球地球的古气候、生物标志化合物以及古海 洋氧化还原条件等研究的深入<sup>[7:9]</sup>以及气候模拟<sup>[10]</sup>,有 些学者认为在冰期期间,水文循环和大陆风化仍然 活跃,地球并未被冰川完全覆盖,尤其是赤道附近存 在一定的开放水域,这就是著名的冰雪地球(hard Snowball Earth)和冰水地球(Slushball Earth)之争<sup>[11]</sup>。目前对 Marinoan 冰期的研究比较深入,包括冰期之间小的冰期—间冰期旋回<sup>[12]</sup>,冰期海洋的氧化还原状态及碳—氮循环过程<sup>[9,13-14]</sup>,生物标志化合物及初级生产力<sup>[15-16]</sup>等,但是对Sturtian 冰期的古海洋环境研究相对较少,尤其缺乏生物地球化学循环的研究。

地质历史时期出现多次无机碳同位素的正偏和 负偏现象,但是目前对其偏移机制仍然存在不同的 认识,可能与成岩作用、生物生产力的增加/降低、有 机质埋藏比例等因素有关<sup>[17-18]</sup>。目前已有很多关于

收稿日期:2023-07-28;修回日期:2023-12-21;录用日期:2024-01-08;网络出版日期:2024-01-08

基金项目:国家自然科学基金项目(42202111);生物地质与环境地质国家重点实验室开放基金(GBL22107);河南省博士后基金(HN2022027);河 南省高等学校重点科研项目(24A170019)[Foundation: National Natural Science Foundation of China, No. 42202111; Open Foundation of State Key Laboratory of Biogeology and Environmental Geology, No. GBL22107; Postdoctoral Foundation of Henan Province, No. HN2022027; Key Scientific Research Project of Colleges and Universities in Henan Province, No. 24A170019] Sturtian冰期之前和冰期之后地层中碳同位素的报 道<sup>[17-18]</sup>,但是对于Sturtian冰期期间碳同位素的组成以 及碳循环过程仍然缺乏系统的研究。南华盆地完整 地记录了南华系的地层序列,包括Sturtian冰期、 Marinoan冰期以及两个冰期之间的间冰期地层,是研 究雪球地球的理想区域。选取南华盆地中黔东松桃 地区高地锰矿区ZK2115钻孔为研究对象,对Sturtian 冰期沉积的铁丝坳组进行详细的有机碳和无机碳同 位素研究,结合地层中的TOC含量,旨在研究Sturtian 冰期期间海洋中的碳同位素组成特征,并探讨极端 气候条件下海洋中的碳循环过程。

## 1 地质背景

中元古代末期一新元古代早期(~1.3~0.9 Ga)形成一个全球性的超大陆一Rodinia超大陆<sup>[19-21]</sup>。在Rodinia超大陆形成过程中,扬子板块与华夏板块碰撞拼合,最终形成华南板块(~820 Ma)<sup>[22]</sup>。新元古代中期,伴随着Rodinia超大陆的裂解,华南板块在~820 Ma开始发生裂谷作用<sup>[23]</sup>,在扬子板块和华夏板块之间形成南华裂谷盆地,在扬子西缘形成康滇裂

谷盆地[23]。

南华裂谷盆地沿北东东向展布<sup>[24-25]</sup>,盆地中发育 武陵次级裂谷盆地和雪峰次级裂谷盆地,二者被天 柱一怀化隆起隔开(图1),在武陵次级裂谷盆地中包 含一系列更次级的地垒和地堑盆地,在这些小的地 堑盆地中发育一系列大型一超大型的锰矿床,被称 为"大塘坡式"锰矿床<sup>[24-25]</sup>。在南华盆地中发育的地 堑盆地完整地记录了南华系的地层序列,地层厚度 变化较大。

研究区位于南华盆地的雪峰次级裂谷盆地,区 内南华纪地层出露较完整,自下而上分别为两界河 组、铁丝坳组、大塘坡组和南沱组。两界河组以中厚 层岩屑砂岩和石英砂岩为主,含白云岩透镜体,地层 厚度变化大,在区域上呈零星分布,主要分布在贵州 东北部的松桃一印江地区,属于陆源碎屑沉积。一 般认为铁丝坳组是Sturtian冰期冰海沉积的产物,主 要由冰碛岩组成,冰碛岩中砾石的分选较差,粒径大 小不一,在0.1 cm×0.2 cm~1 cm×3 cm,磨圆也较差, 并且砾石的含量自下而上逐渐变少。铁丝坳组与下 伏两界河组整合接触。大塘坡组代表Sturtian冰期



Fig.1 (a) Structure of the Neoproterozoic Nanhua rift basin, located in southeastern margin of the Yangtze Block, South China<sup>[25]</sup>;
 (b) geological map of Wuling secondary rift basin during the early Nanhua period in the Guizhou-Hunan-Chongqing adjacent area<sup>[25]</sup>
 I-1. Wuling secondary rift basin; I-2. Tianzhu-Huaihua uplift; I-3. Xuefeng secondary rift basin

岩组成,页岩底部包含一套富锰页岩层,是"大塘坡 式"锰矿的富集层位,上部主要由灰绿色粉砂岩组 成。锰矿层中的锰主要以菱锰矿(MnCO<sub>3</sub>)的形式存 在,从矿床的中心到边缘,矿床品位逐渐降低,块状 矿石逐渐变为条带状矿石。大塘坡组与下伏铁丝坳 组和上覆南沱组呈整合接触。南沱组代表Marinoan 冰期期间的冰海沉积,主要由砂泥质冰碛岩组成。 选取黔东松桃高地锰矿区ZK2115钻孔为研究对象, 该钻孔到铁丝坳组底部终止,与大塘坡组(约231 m) 和南沱组(约284 m)相比,铁丝坳组较薄(约10 m) (图2)。ZK2115钻孔铁丝坳组冰碛岩中可见大小不 等的砾石,并且在冰碛岩基质中发现很多细小的岩 石碎屑,例如石英、长石等(图3)。

Sturtian冰期的开始和结束在全球范围内是同步 的<sup>[3,26]</sup>,精确的年代学研究将Sturtian冰期的初始时间 限定在717 Ma左右<sup>[1-2,27]</sup>。目前普遍认为Sturtian冰期 包含两幕冰期事件,在华南地区分别对应年龄较老 的长安冰期和较新的古城冰期[28],并且越来越多的研 究表明 Sturtian 冰期的第二幕冰期在全球范围内也 是同步的,其开始的年龄在690 Ma左右<sup>[28-29]</sup>,结束的 年龄对应整个Sturtian冰期结束的年龄,即660 Ma左 右[30-33]。在扬子东南缘南华裂谷盆地中两界河组底 部最年轻的单颗粒锆石U-Pb年龄为708±15 Ma<sup>[34]</sup>,区 域上相当层位的渫水河组顶部凝灰质粉砂岩中的锆 石 SIMS U-Pb 年龄为 691.9±8.0 Ma<sup>[28]</sup>。这些年代学资 料限定两界河组的沉积时代介于708~690 Ma。除此 之外,两界河组的化学风化指标——CIA(Chemical Index of Alteration)整体上介于65~75<sup>[35]</sup>,代表了当时 温暖湿润的古气候,同时也说明两界河组属于 Sturtian冰期两次幕式冰期之间的间冰期沉积,铁丝 坳组代表了 Sturtian 冰期的第二幕冰期沉积。





## 2 样品测试和结果

研究的样品均来自黔东松桃地区高地锰矿区 ZK2115钻孔,该钻孔中铁丝坳组地层厚度约为10m, 采集样品的间隔为0.3~0.6m,共采集铁丝坳组样品 24件。由于铁丝坳组样品主要为Sturtian冰期期间 沉积的冰碛岩,岩石中含有大小不等的砾石,因此将 固体的岩石样品研磨成粉末(200目)之前,需要剔除 掉样品中的砾石,地球化学测试的对象是冰碛岩的 基质部分。



图 3 铁丝坳组冰碛岩手标本(a)和基质部分偏光显微镜(b)照片 Fig.3 Rock photo (a) and matrix microscope photo (b) of diamictites in the Tiesi'ao Formation

 $δ^{13}C_{catb} 和 \delta^{18}O 的测试: 首先称量适量样品置于反应瓶中, 连接至真空系统, 然后与高纯磷酸在 70°C反应, 将收集到的 CO<sub>2</sub>进行 C-O 同位素测试, 该过程在 kiel IV-MAT253 连用系统上进行, 采用的国际标样为 GBW04416 (<math>\delta^{13}C=+1.61\%$ ,  $\delta^{18}O=-11.59\%$ )和 GBW04417( $\delta^{13}C=-6.06\%$ ,  $\delta^{18}O=-24.12\%$ ), 分析结果以 Vienna Pee Dee Belemnite(V-PDB)为标准, 分析精度小于 0.1‰。

在测试δ<sup>13</sup>C<sub>org</sub>之前需要去除掉样品中的无机碳部 分,具体方法如下:首先取3~5g样品放到50mL离心 管中,然后少量多次加入4mol/L的盐酸,搅拌均匀, 并静置2h,然后用离心机离心,去除上清液,重复上 述过程2~3次,确保除掉样品中的无机碳被完全去 除。向残余物中加入超纯水,摇晃均匀,然后离心, 去掉上清液,重复上述过程5~6次,直至离心之后的 上清液呈中性。最后将水洗之后的残余物放在 65°C的烘箱中,待样品烘干后磨成粉末备用。称量 适量样品,用锡纸杯包裹紧密之后在EA+MAT253上 进行测试。采用的国际标样为GBW04407(-22.43±0.07‰)和GBW04408(-36.91±0.10‰),国内标样ACET(-26.33‰)分析结果以V-PDB为标准,分析精度小于0.06‰。

研究中涉及的无机碳一氧同位素和有机碳同位 素的测试均在中国地质大学(武汉)地质过程与矿产 资源国家重点实验室完成,测试结果如表1所示。

铁丝坳组 $\delta^{13}C_{carb}$ 介于-9.29‰~-3.37‰,平均值为-7.24‰, $\delta^{18}$ 〇介于-15.91‰~-7.37‰,平均值为-12.71‰; $\delta^{13}C_{org}$ 介于-33.36‰~-23.35‰,平均值为-29.29‰。此外,铁丝坳组的碳同位素分馏( $\Delta^{13}C$ ; $\delta^{13}C_{carb}$ - $\delta^{13}C_{org}$ )介于17.84‰~25.90‰。研究样品整体上TOC含量相对较低(0.2±0.6%),但是在铁丝坳组顶部个别样品的TOC含量相对较高,最高可达2.7%<sup>[36]</sup>(图4)。

在本研究的测试数据中,δ<sup>13</sup>C<sub>eatb</sub>和δ<sup>13</sup>C<sub>org</sub>从铁丝坳 组底部开始存在一个明显的上升趋势,随后又呈下 降的趋势,Δ<sup>13</sup>C从铁丝坳组的底部到顶部具有明显

表1 黔东松桃地区高地锰矿区 ZK2115 钻孔铁丝坳组地球化学测试结果

Table 1 Geochemical results of the Tiesi'ao Formation in drillcore ZK2115, Gaodi manganese deposit,

Songtao area, eastern Guizhou province

样品编号	深度/m	$\delta^{13}\mathrm{C_{carb}}$ /%o	$\delta^{ m ^{18}O}$ /% $o$	$\delta^{13} \mathrm{C_{org}} / \% o$	$\triangle^{13}C_{carb-org}/\%$	TOC/%
ZK2115-H51	1 637.98			-30.22		0.2
ZK2115-H52	1 638.28			-29.50		0.1
ZK2115-H53	1 638.58	-7.45	-14.06	-33.18	25.73	0.7
ZK2115-H54	1 638.88	-7.31	-15.89	-33.21	25.90	0.7
ZK2115-H55	1 639.18			-33.36		2.7
ZK2115-H56	1 639.48	-8.05	-11.78	-30.49	22.44	0.1
ZK2115-H57	1 639.78	-8.21	-13.58	-30.86	22.65	0.1
ZK2115-H58	1 640.08	-8.18	-11.72	-30.17	21.99	0.1
ZK2115-H59	1 640.38			-29.93		0.1
ZK2115-H61	1 640.98	-8.16	-10.53	-30.07	21.91	0.1
ZK2115-H63	1 641.58	-9.29	-11.91	-30.30	21.01	0.1
ZK2115-H64	1 641.88			-30.07		0.1
ZK2115-H65	1 642.48			-29.40		0.1
ZK2115-H66	1 642.78			-27.95		0.1
ZK2115-H68	1 643.38	-4.91	-7.84	-26.87	21.96	0
ZK2115-H69	1 643.68			-29.00		0.1
ZK2115-H71	1 644.28	-6.61	-15.14	-27.57	20.96	0.1
ZK2115-H72	1 644.58	-6.80	-14.86	-27.15	20.35	0
ZK2115-H74	1 645.18	-3.37	-7.37	-23.35	19.98	0.1
ZK2115-H75	1 645.48	-7.11	-11.42	-27.89	20.78	0.1
ZK2115-H77	1 646.08	-6.91	-14.14	-27.43	20.52	0.1
ZK2115-H78	1 646.38	-7.43	-14.57	-27.01	19.58	0.1
ZK2115-H79	1 646.68	-8.86	-15.91	-26.70	17.84	0
ZK2115-H82	1 647.58			-31.21		0.1

注:TOC数据来自文献[36]。



图 4 黔东松桃高地锰矿区 ZK2115 钻孔铁丝坳组 $\delta^{13}C_{carb}, \delta^{18}O, \delta^{13}C_{org}, \Delta^{13}C \, 以及 TOC 分布情况(TOC 引自文献[36])$ Fig.4 The  $\delta^{13}C_{carb}, \delta^{18}O, \delta^{13}C_{org}, \Delta^{13}C$  and TOC profiles of the Tiesi 'ao Formation in drillcore ZK2115, Gaodi manganese deposit, Songtao area, eastern Guizhou province (TOC data are from reference [36])

的上升趋势(图4)。 $\delta^{13}C_{earb}$ 和 $\delta^{18}O$ 显示正相关关系(r=+0.55,  $p(\alpha)<0.02$ , n=15;图5a),  $\delta^{13}C_{earb}$ 和 $\delta^{13}C_{org}$ 也显示 正相关关系(r=+0.61,  $p(\alpha)<0.01$ , n=15;图5b),  $\delta^{13}C_{earb}$ 和TOC没有相关性(r=-0.12,  $p(\alpha)<0.1$ , n=15;图5c),  $\delta^{13}C_{org}$ 和TOC显示微弱的负相关性(r=-0.55,  $p(\alpha)<0.01$ , n=24;图5d)。 $\delta^{13}C_{earb}$ 和 $\Delta^{13}C$ 不显示相关性(r=-0.06,  $p(\alpha)<0.1$ , n=15;图5e), 但是 $\delta^{13}C_{org}$ 和 $\Delta^{13}C$ 显示 很好的负相关关系(r=-0.83,  $p(\alpha)<0.001$ , n=15; 图5f)。

## 3 讨论

## 3.1 碳同位素数据的可用性分析

沉积碳酸盐岩记录了当时的古海洋环境信息, 但是碳酸盐岩在后期成岩过程中容易受到成岩作用 的影响,导致其微量元素(Mn、Fe、Ca和Sr)含量以及 δ<sup>13</sup>C<sub>catb</sub>和δ<sup>18</sup>O值的变化<sup>[37-38]</sup>。因此,在利用碳酸盐岩进 行古海洋环境研究之前,需要考虑成岩作用是否影 响碳酸盐岩中记录的古海水初始信号。

在成岩过程中,沉积碳酸盐岩晶格中的Ca和Sr 会被Fe和Mn取代,导致Fe和Mn含量的增加,Ca和 Sr含量的降低,因此Mn/Sr、Fe/Sr比值可以作为沉积 碳酸盐岩成岩蚀变的指标<sup>[37,39]</sup>。一般认为Mn/Sr<2, Fe/Sr<50时,碳酸盐岩仍然保存了原始的碳同位素组 成特征<sup>[39-40]</sup>,但是在前寒武纪时期,原始沉积的碳酸 盐岩矿物中铁、锰含量本来就比较高,并不是后期成 岩作用导致的<sup>[41]</sup>。因此,对于本研究中的样品,Mn/Sr 和Fe/Sr不适合作为成岩作用的指标。

成岩作用会导致碳酸盐岩δ<sup>13</sup>C<sub>carb</sub>和δ<sup>18</sup>O的降低, 并且δ<sup>18</sup>O比δ<sup>13</sup>C<sub>carb</sub>更容易受到成岩作用的影响<sup>137,39-40</sup>]。 碳酸盐岩δ<sup>18</sup>O值可以作为成岩作用的判别指标<sup>[40]</sup>,一 般认为当δ<sup>18</sup>O>-10‰ 时,碳酸盐岩受到的成岩作用 比较微弱<sup>[39]</sup>。此外,δ<sup>13</sup>C<sub>carb</sub>和δ<sup>18</sup>O的相关性也被用来 评估成岩作用对碳酸盐岩的影响<sup>[37-39]</sup>,当成岩作用较 强烈时,会同时降低δ<sup>13</sup>C<sub>carb</sub>和δ<sup>18</sup>O的值,δ<sup>13</sup>C<sub>carb</sub>和δ<sup>18</sup>O 显示正相关关系。当地层中出现耦合的δ<sup>13</sup>C<sub>carb</sub>和δ<sup>18</sup>O 显示正相关关系。当地层中出现耦合的δ<sup>13</sup>C<sub>carb</sub>和δ<sup>18</sup>O 显示正相关关系。当地层中出现耦合的δ<sup>13</sup>C<sub>carb</sub>和δ<sup>18</sup>O 显示正相关关系。当地层中出现耦合的δ<sup>13</sup>C<sub>carb</sub>和δ<sup>18</sup>O 显示正相关关系。当地层中出现耦合的δ<sup>13</sup>C<sub>carb</sub>和δ<sup>18</sup>O

南华盆地 Sturtian 冰期期间沉积的铁丝坳组 样品中 $\delta^{18}O$ 值介于-15.91%~-7.37%(平均值为 -12.71%),大部分样品的 $\delta^{18}O$ 都小于-10%,并且  $\delta^{13}C_{earb}-\delta^{18}O$ 呈现中等程度的正相关关系(图 5a),似乎 预示着成岩作用对初始古海洋信号的改变。但是铁 丝坳组冰碛岩基质 $\delta^{13}C_{earb}$ (-9.29‰~-3.37‰,平均值



图 5 ZK2115 钻孔铁丝坳组<sup>13</sup>C<sub>aub</sub>-<sup>18</sup>O, <sup>13</sup>C<sub>aub</sub>-<sup>13</sup>C<sub>aub</sub>, <sup>31</sup>C<sub>aub</sub>-<sup>7</sup>TOC, δ<sup>13</sup>C<sub>aub</sub>-<sup>7</sup>TOC, δ<sup>13</sup>C<sub>aub</sub>-<sup>Δ</sup><sup>13</sup>C, δ<sup>13</sup>C<sub>aub</sub>-<sup>Δ</sup><sup>13</sup>C, δ<sup>13</sup>C<sub>aub</sub>-<sup>Δ</sup><sup>13</sup>C, δ<sup>13</sup>C<sub>aub</sub>-<sup>Δ</sup><sup>13</sup>C, δ<sup>13</sup>C<sub>aub</sub>-<sup>Δ</sup><sup>13</sup>C, δ<sup>13</sup>C, δ<sup>13</sup>C Fig.5 Crossplots of  ${}^{13}C_{earb} - {}^{18}O, {}^{13}C_{earb} - {}^{13}C_{org}, \delta {}^{13}C_{earb} - TOC, \delta {}^{13}C_{org} - TOC, \delta {}^{13}C_{earb} - \Delta {}^{13}C, \delta {}^{13}C_{org} - \Delta {}^{13}C$  in the Tiesi 'ao Formation, drillcore ZK2115

-2

15 L -35

-30

为-7.24‰)和δ<sup>13</sup>C<sub>org</sub>(-33.36‰~-23.35‰,平均值为 -29.29‰)显示明显的正相关关系(图5b),这与成岩 作用相悖,因为在前寒武纪时期没有任何一种机制 可能同时改变 $\delta^{13}C_{carb}$ 和 $\delta^{13}C_{org}$ 值<sup>[43]</sup>。因此,我们认为铁 丝坳组记录了原始的碳同位素组成信号,这与地质 历史时期其他地层记录一致<sup>[42-43]</sup>,但是δ<sup>18</sup>O可能受到 了成岩作用的影响。

-8

-6

 $\delta^{13}C_{carb}/\% o$ 

-4

#### 3.2 Sturtian 冰期期间的光合作用

在新元古代雪球地球期间,最开始认为该时期 整个地球几乎完全被冰雪覆盖,陆地、海洋和大气之 间的物质循环受到阻碍,生态系统崩塌,光合作用停 滞,海洋处于缺氧状态<sup>[46]</sup>,并且认为海洋中的DIC库 δ<sup>13</sup>C 值在-5‰ 左右<sup>[5]</sup>, 与地幔δ<sup>13</sup>C 值接近<sup>[44]</sup>。随着对 冰期沉积地层越来越深入的研究以及气候模型模 拟,一些学者对雪球地球的冰川范围产生了不同的 意见,认为在冰期期间,地球并未完全被冰川覆盖, 部分地区仍然存在开放水域[9-10]。化学风化指标 (CIA)表明冰期期间化学风化作用仍在进行<sup>[8,45]</sup>,并且 冰期期间水文循环并未停止[45],陆地和海洋之间仍然 存在物质交换。

-25

 $\delta^{13}C_{arg}/\%$ 

-20

在雪球地球期间,全球生态系统并未完全崩溃, 生命主要以微观和软体的形式存在1460。有机分子证 据以及生物标志化合物的证据表明,在冰期期间存 在微弱的透光带,光合作用比较微弱,虽然生物生产 力很低,但是从未停止,并且沉积有机质主要来源于 海洋中的光合自养生物[47-48]。在澳大利亚和斯瓦尔 巴群岛 Sturtian 冰期沉积的冰碛岩中已经发现有机 壁微化石,尽管其多样性很低,但是其分布广泛且相

当丰富<sup>[49]</sup>。在巴西东南部 Sturtian 冰期期间沉积地 层中发育横向上广泛分布的黑色页岩层(TOC可 达 3.0%), 远远高于上下冰碛岩层的 TOC 含量 (< 0.5%), 地层中提取出的生物标志化合物表明 Sturtian冰期期间,该地区存在一个复杂而多产的微 生物生态系统,包括光合细菌和真核生物,光合作用 仍在进行,表层海水处于氧化状态,当时海洋中可能 存在很薄的冰或者处于无冰状态。阿曼地区 Marinaon冰期沉积物中发现了24-异丙基胆甾烷,该 物质被认为是海洋中海绵动物产生的Cau甾醇的碳氢 化合物残留物,代表了化石记录中最古老的动物证 据<sup>[15]</sup>;华南 Marinoan 冰期冰碛岩中的页岩层中首次发 现了底栖大型藻类<sup>116]</sup>。这些化石数据都表明真核生 物在 Marinaon 冰期幸存下来,并未由于冰期恶劣的 环境而消失。纳米比亚 Marinoan 冰期沉积的 Ghaub 组和同时期的华南南沱组的铁组分以及C-N同位素 等地球化学数据的研究表明,冰期期间存在显著的 开阔水域,生物地球化学循环较为活跃,生物生产力 以及产氧作用存在,并且表层海水含氧,深部水体 缺氧[9,14]。

在新元古代南华盆地中, Sturtian 冰期期间沉积 的铁丝坳组TOC含量很低,除了一个样品(黑色页 岩)达到2.7%,其余样品都小于1.0%。铁丝坳组的 低 TOC 含量也与其他地区 Sturtian 冰期期间沉积冰 碛岩中的TOC含量一致,例如华南地区江口组[48]和 澳大利亚Wilyerpa组<sup>[50]</sup>。Marinaon冰期期间,纳米比 亚地区沉积的 Ghaub 组<sup>144</sup>和华南地区沉积的南沱 组<sup>[9,47-48,51]</sup>冰碛岩中TOC含量也很低(<0.2%),与 Sturtian冰期类似。沉积物中有机质的来源一般分为 两种,一种是光合作用产生的有机质,另外一种是陆 源输入的有机质。铁丝坳组δ<sup>13</sup>C<sub>ent</sub>和δ<sup>13</sup>C<sub>ort</sub>具有明显 的正相关关系(图5b),表明地层中的有机质主要是 水体中光合作用的产物,有机质中的碳来源于海洋 中的DIC库。以上证据均表明Sturtian冰期期间的生 态系统仍然存在,光合作用仍在进行。有机质的矿 化作用是影响地层中有机质保存的重要因素,在新 元古代海水中硫酸盐的浓度很低521,通过硫酸盐还原 等过程消耗的有机质的量特别少,有机质矿化作用 比较微弱,地层中TOC的含量主要受到光合作用产 生的有机质的影响。Sturtian冰川沉积物中TOC的含 量比间冰期低1~2个数量级[36.47-48],尽管冰川沉积物 的沉积速率相对较高,但是仍然可以说明冰室气候 条件下光合作用的速率比较缓慢,初级生产力水平 较低,通过光合作用产生的有机质十分有限。

#### 3.3 Sturtian冰期碳同位素的负偏移

黔东松桃地区高地ZK2115钻孔铁丝坳组冰 碛岩基质部分δ<sup>13</sup>C<sub>eat</sub>值介于-9.29‰~-3.37‰(平均值 为-7.24‰),与研究区临近的ZK405钻孔同一层位冰 碛岩基质的δ<sup>13</sup>C<sub>eat</sub>值(-11.3% 到-8.3%,平均值为 -9.6‰)<sup>[53]</sup>一致,存在明显的碳同位素负偏。铁丝坳 组无机碳同位素代表原始的碳同位素组成信号,并 且在冰期沉积物沉积过程中矿化的有机质很少,因 此该过程产生的无机碳对海洋 DIC 库的影响很小, 铁丝坳组δ<sup>13</sup>C<sub>ent</sub>可以代表当时水体中DIC库的碳同 位素组成。桂西地区Sturtian冰期沉积的富禄组发 育一套碳酸盐岩夹层,可能代表了冰期期间一次小 的间冰期沉积,碳酸盐岩夹层 $\delta^{13}C_{aut}$ 介于-5.0‰~ -3.0‰(平均值为-3.5‰)<sup>[54]</sup>,与Sturtian之后的盖帽白 云岩类似<sup>[30]</sup>,并且δ<sup>13</sup>C<sub>eat</sub>与δ<sup>13</sup>C<sub>eat</sub>(-27.8%--22.2%)呈 现明显的解耦现象[54],这可能与环境变化导致的碳循 环扰动有关。此外,世界其他地区也报道了Sturtian 冰期沉积的冰碛岩中无机碳同位素数据。例如,在 澳大利亚 Sturtian 冰期沉积的 Wilverpa 组,沉积过程 中形成的白云石δ<sup>13</sup>C<sub>ent</sub>介于-5.3‰~+2.0‰(平均值为 -1.7‰)<sup>[50,55]</sup>;在斯瓦尔巴特群岛, Sturtian冰期沉积的 Petrovbreen 段冰碛岩基质δ<sup>13</sup>C<sub>cath</sub>介于-5.0‰~+2.0‰ [56];在纳米比亚,Sturtian冰期沉积的Chuos组,其泥晶 白云岩和灰岩层δ<sup>13</sup>C<sub>erb</sub>介于-1.9%~-9.2% (平均值为 -7.3‰)[57]。这些证据表明 Sturtian 冰期期间,海洋中 的DIC库存在负偏,其 $\delta^{13}$ C值低于现代海洋的 $\delta^{13}$ C。 (0左右)。此外,南华盆地 Sturtian 冰川沉积物中 δ<sup>13</sup>C<sub>eath</sub>值低于 Sturtian 冰期之前(0~-5.0‰)<sup>[56-57]</sup>和冰期 之后(-4.0%~+4.0%)[30.57]的地层(图6)。南华盆地在 Sturtian冰期期间海水 DIC 库可能受到大气 CO<sub>2</sub>的溶 解,火山/热液活动的输入以及有机质矿化的影响,但 是大气中CO,在水中溶解的 $\delta^{13}C_{ext}$ (-4.0%~ -5.0‰)<sup>[5]</sup>,火山或者热液活动δ<sup>13</sup>C<sub>art</sub> (-5.0‰~ -7.0‰)<sup>[5]</sup>均高于水体中DIC库的δ<sup>13</sup>C<sub>aut</sub>,并且冰期期 间有机质矿化作用较微弱,可以忽略,因此冰期DIC 库碳同位素的负偏需要其他<sup>13</sup>C缺乏的碳源的输入。 Sturtian 冰期期间 $\delta^{13}C_{ext}$ 的负偏可能与DIC 库碳同位 素的大规模扰动有关,受到环境因素的制约,也可能 受到全球碳循环过程的影响[43.58],对于其负偏机制目 前仍然无法确定,缺乏实质性的证据。





#### 3.4 Sturtian冰期碳循环及对海洋状态的启示

Sturtian 冰期沉积的铁丝坳组δ<sup>13</sup>C<sub>eat</sub>代表了当时 水体 DIC 库的碳同位素组成信息,在-7‰左右,具有 明显的负偏。在冰期期间,全球处于冰室气候状态, 生态系统并未完全崩塌,光合作用仍在进行,但是光 合作用的速率极低。此外,有机质的矿化作用比较微 弱,光合作用产生的有机质大部分都在地层中保存下 来,由于初级生产力很低,产生的有机质有限,在地层 中保存下来的TOC 含量很低(0.2%)。随着冰室气候 逐渐向温室气候转变<sup>18</sup>,温度的升高可能导致有机质 和 DIC 库之间的碳同位素分馏逐渐增大,△<sup>13</sup>C 值从 铁丝坳组底部到顶部具有明显的上升趋势。

在 Sturtian 冰期期间,有机碳和无机碳之间存在 比较活跃的碳循环过程。此外,铁丝坳组古气候指 标——CIA 介于 45~67(平均值为 58),并且从铁丝坳 组的底部到顶部整体上显示上升的趋势,表明在冰 室气候环境下,化学风化作用并未完全停止<sup>[8]</sup>。铁丝 坳组  $\delta^{13}C_{eat}$ 的升高—降低趋势也反映了当时的水体 不是封闭的环境,而是开放的环境,海洋中的碳循环 受到扰动。如果是封闭的环境,那么随着海水中的 无机碳通过光合作用固定在有机质中,那么水体中 将会越来越富集<sup>13</sup>C,导致 DIC 库的  $\delta^{13}C_{eat}$ 值逐渐升 高,这与实际情况相悖。以上证据均支持"冰水地 球"假说<sup>[10]</sup>,认为在 Sturtian 冰期这种极端气候条件 下,部分地区仍然存在开放水域,陆地和海洋之间的 物质交换仍然存在,生物地球化学循环仍然比较 活跃。

## 4 结论

(1) Sturtian冰期期间,南华盆地中的DIC库明显 缺乏<sup>13</sup>C,具有较低的δ<sup>13</sup>C值(-7‰左右),铁丝坳组具 有明显的δ<sup>13</sup>C<sub>eat</sub>负偏现象。

(2) Sturtian 冰期期间,虽然光合作用在极端条件下仍在进行,但光合速率极低,铁丝坳组中保存下来的有机质主要是通过光合作用产生的,并且有机质中的碳主要来源于海洋中的DIC库。

(3) Sturtian 冰期期间,地球并未被冰川完全覆盖,存在开放水域,生物地球化学循环过程仍然比较活跃。

致谢 感谢中国地质大学(武汉)生物地质与 环境地质国家重点实验室常标副研究员和薛书雨博 士在实验室测试过程中提供的帮助。感谢审稿专家 在本文修改过程中提出的宝贵意见。

## 参考文献(References)

- Macdonald F A, Schmitz M D, Crowley J L, et al. Calibrating the Cryogenian[J]. Science, 2010, 327(5970): 1241-1243.
- [2] Lan Z W, Li X H, Zhu M Y, et al. A rapid and synchronous initiation of the wide spread Cryogenian glaciations[J]. Precambrian Research, 2014, 255: 401-411.
- [3] Rooney A D, Strauss J V, Brandon A D, et al. A Cryogenian chronology: Two long-lasting synchronous Neoproterozoic glaciations
   [J]. Geology, 2015, 43(5): 459-462.
- [4] Kirschvink J L. Late Proterozoic low-latitude global glaciation: The Snowball Earth[M]//Schopf J W, Klein C. The Proterozoic

biosphere: A multidisciplinary study. New York: Cambridge University Press, 1992: 51-52.

- [5] Hoffman P F, Kaufman A J, Halverson G P, et al. A Neoproterozoic Snowball Earth[J]. Science, 1998, 281(5381): 1342-1346.
- [6] Hoffman P F, Schrag D P. The Snowball Earth hypothesis: Testing the limits of global change[J]. Terra Nova, 2002, 14(3): 129-155.
- [7] Olcott A N, Sessions A L, Corsetti F A, et al. Biomarker evidence for photosynthesis during Neoproterozoic glaciation[J]. Science, 2005, 310(5747): 471-474.
- [8] Wang P, Du Y S, Yu W C, et al. The chemical index of alteration (CIA) as a proxy for climate change during glacial-interglacial transitions in Earth history[J]. Earth-Science Reviews, 2020, 201: 103032.
- [9] Song H Y, An Z H, Ye Q, et al. Mid-latitudinal habitable environment for marine eukaryotes during the waning stage of the Marinoan snowball glaciation[J]. Nature Communications, 2023, 14 (1): 1564.
- [10] Hyde W T, Crowley T J, Baum S K, et al. Neoproterozoic 'Snowball Earth' simulations with a coupled climate/ice-sheet model[J]. Nature, 2000, 405(6785): 425-429.
- [11] Micheels A, Montenari M. A Snowball Earth versus a slushball Earth: Results from Neoproterozoic climate modeling sensitivity experiments[J]. Geosphere, 2008, 4(2): 401-410.
- [12] Lang X G, Chen J T, Cui H, et al. Cyclic cold climate during the Nantuo Glaciation: Evidence from the Cryogenian Nantuo Formation in the Yangtze Block, South China[J]. Precambrian Research, 2018, 310: 243-255.
- [13] 沈洪娟,顾尚义,赵思凡,等.华南南华纪南沱冰期海洋环境的沉积地球化学记录:来自黔东部南华系南沱组白云岩碳氧同位素和微量元素的证据[J].地质论评,2020,66(1):214-228. [Shen Hongjuan, Gu Shangyi, Zhao Sifan, et al. The sedimentary geochemical records of ocean environment during the Nantuo (Marinoan) glaciation in South China: Carbon and oxygen isotopes and trace element compositions of dolostone in Nantuo Formation, Nanhuan System, in eastern Guizhou[J]. Geological Review, 2020, 66(1): 214-228. ]
- [14] Johnson B W, Poulton S W, Goldblatt C. Marine oxygen production and open water supported an active nitrogen cycle during the Marinoan Snowball Earth[J]. Nature Communications, 2017, 8(1): 1316.
- [15] Love G D, Grosjean E, Stalvies C, et al. Fossil steroids record the appearance of Demospongiae during the Cryogenian period
   [J]. Nature, 2009, 457(7230): 718-721.
- [16] Ye Q, Tong J N, Xiao S H, et al. The survival of benthic macroscopic phototrophs on a Neoproterozoic Snowball Earth[J]. Geology, 2015, 43(6): 507-510.
- [17] Cox G M, Halverson G P, Stevenson R K, et al. Continental flood basalt weathering as a trigger for Neoproterozoic Snowball Earth[J]. Earth and Planetary Science Letters, 2016, 446: 89-99.

- [18] Ward J F, Verdel C, Campbell M J, et al. Rare earth element geochemistry of Australian Neoproterozoic carbonate: Constraints on the Neoproterozoic oxygenation events[J]. Precambrian Research, 2019, 335: 105471.
- [19] Dalziel I W D. Pacific margins of Laurentia and East Antarctica-Australia as a conjugate rift pair: Evidence and implications for an Eocambrian supercontinent[J]. Geology, 1991, 19(6): 598-601.
- [20] Hoffman P F. Did the breakout of Laurentia turn Gondwanaland inside-out?[J]. Science, 1991, 252(5011): 1409-1412.
- [21] Moores E M. Southwest U. S. -East Antarctic (SWEAT) connection: A hypothesis[J]. Geology, 1991, 19(5): 425-428.
- [22] Li X H, Li W X, Li Z X, et al. Amalgamation between the Yangtze and Cathaysia Blocks in South China: Constraints from SHRIMP U-Pb zircon ages, geochemistry and Nd-Hf isotopes of the Shuangxiwu volcanic rocks[J]. Precambrian Research, 2009, 174(1/2): 117-128.
- [23] Wang J, Li Z X. History of Neoproterozoic rift basins in South China: Implications for Rodinia break-up[J]. Precambrian Research, 2003, 122(1/2/3/4): 141-158.
- [24] 杜远生,周琦,余文超,等. Rodinia超大陆裂解、Sturtian冰期 事件和扬子地块东南缘大规模锰成矿作用[J]. 地质科技情报, 2015,34(6):1-7. [Du Yuansheng, Zhou Qi, Yu Wenchao, et al. Linking the Cryogenian manganese metallogenic process in the southeast margin of Yangtze Block to break-up of Rodinia supercontinent and Sturtian Glaciation[J]. Geological Science and Technology Information, 2015, 34(6): 1-7. ]
- [25] 周琦,杜远生,袁良军,等. 黔湘渝毗邻区南华纪武陵裂谷盆 地结构及其对锰矿的控制作用[J]. 地球科学,2016,41(2): 177-188. [Zhou Qi, Du Yuansheng, Yuan Liangjun, et al. The structure of the Wuling rift basin and its control on the manganese deposit during the Nanhua period in Guizhou-Hunan-Chongqing border area, South China[J]. Earth Science, 2016, 41 (2): 177-188. ]
- [26] Hoffman P F, Abbot D S, Ashkenazy Y, et al. Snowball Earth climate dynamics and Cryogenian geology-geobiology[J]. Science Advances, 2017, 3(11): e1600983.
- [27] Lan Z W, Huyskens M H, Lu K, et al. Toward refining the onset age of Sturtian Glaciation in South China[J]. Precambrian Research, 2020, 338: 105555.
- [28] Lan Z W, Li X H, Zhang Q R, et al. Global synchronous initiation of the 2nd episode of Sturtian glaciation: SIMS zircon U-Pb and O isotope evidence from the Jiangkou Group, South China [J]. Precambrian Research, 2015, 267: 28-38.
- [29] Balgord E A, Yonkee W A, Link P K, et al. Stratigraphic, geochronologic, and geochemical record of the Cryogenian Perry Canyon Formation, northern Utah: Implications for Rodinia rifting and Snowball Earth glaciation[J]. GSA Bulletin, 2013, 125 (9/10): 1442-1467.
- [30] Yu W C, Algeo T J, Du Y S, et al. Newly discovered Sturtian

cap carbonate in the Nanhua Basin, South China[J]. Precambrian Research, 2017, 293: 112-130.

- [31] Zhou C M, Tucker R, Xiao S H, et al. New constraints on the ages of Neoproterozoic glaciations in South China[J]. Geology, 2004, 32(5): 437-440.
- [32] Zhou C M, Huyskens M H, Lang X G, et al. Calibrating the terminations of Cryogenian global glaciations[J]. Geology, 2019, 47(3): 251-254.
- [33] Rooney A D, Yang C, Condon D J, et al. U-Pb and Re-Os geochronology tracks stratigraphic condensation in the Sturtian Snowball Earth aftermath[J]. Geology, 2020, 48(6): 625-629.
- [34] 高永娟,张予杰,安显银,等. 扬子东南缘两界河组碎屑锆石 U-Pb年龄及其对 Sturtian冰川作用的启示[J]. 地球科学, 2020,45(8):3070-3081. [Gao Yongjuan, Zhang Yujie, An Xianyin, et al. Detrital zircon U-Pb ages of Liangjiehe Formation in east Guizhou province and its implications for Sturtian Glaciation [J]. Earth Science, 2020, 45(8): 3070-3081.]
- [35] 郑杰,阳正熙,刘石磊,等. 黔东北地区南华系两界河组CIA 指数特征及意义[J]. 沉积与特提斯地质,2019,39(1):50-59.
  [Zheng Jie, Yang Zhengxi, Liu Shilei, et al. The chemical alteration indexes and their significance for the Nanhuan Liangjiehe Formation in northeastern Guizhou[J]. Sedimentary Geology and Tethyan Geology, 2019, 39(1): 50-59.
- [36] Wang P, Algeo T J, Zhou Q, et al. Large accumulations of <sup>34</sup>Senriched pyrite in a low-sulfate marine basin: The Sturtian Nanhua Basin, South China[J]. Precambrian Research, 2019, 335: 105504.
- [37] Swart P K. The geochemistry of carbonate diagenesis: The past, present and future[J]. Sedimentology, 2015, 62(5): 1233-1304.
- [38] Swart P K, Oehlert A M. Revised interpretations of stable C and O patterns in carbonate rocks resulting from meteoric diagenesis[J]. Sedimentary Geology, 2018, 364: 14-23.
- [39] Kaufman A J, Knoll A H. Neoproterozoic variations in the Cisotopic composition of seawater: Stratigraphic and biogeochemical implications[J]. Precambrian Research, 1995, 73(1/2/3/4): 27-49.
- [40] Jacobsen S B, Kaufman A J. The Sr, C and O isotopic evolution of Neoproterozoic seawater[J]. Chemical Geology, 1999, 161(1/ 2/3): 37-57.
- [41] Yoshioka H, Asahara Y, Tojo B, et al. Systematic variations in C, O, and Sr isotopes and elemental concentrations in Neoproterozoic carbonates in Namibia: Implications for a glacial to interglacial transition[J]. Precambrian Research, 2003, 124(1): 69-85.
- [42] Oehlert A M, Swart P K. Interpreting carbonate and organic carbon isotope covariance in the sedimentary record[J]. Nature Communications, 2014, 5(1): 4672.
- [43] Johnston D T, Macdonald F A, Gill B C, et al. Uncovering the Neoproterozoic carbon cycle[J]. Nature, 2012, 483(7389): 320-323.
- [44] Kump L R. Interpreting carbon-isotope excursions: Strangelove

oceans[J]. Geology, 1991, 19(4): 299-302.

- [45] Rieu R, Allen P A, Plötze M, et al. Climatic cycles during a Neoproterozoic "Snowball" glacial epoch[J]. Geology, 2007, 35(4): 299-302.
- [46] Corsetti F A, Olcott A N, Bakermans C. The biotic response to Neoproterozoic Snowball Earth[J]. Palaeogeography, Palaeoclimatology, Palaeoecology, 2006, 232(2/3/4): 114-130.
- [47] 李美俊,王铁冠. 扬子区新元古代"雪球"时期古环境的分子 地球化学证据[J]. 地质学报,2007,81(2):220-229. [Li Meijun, Wang Tieguan. Molecular geochemical evidence for the paleoenvironment of the Late Neoproterozoic "Snowball Earth" age in the Yangtze region[J]. Acta Geologica Sinica, 2007, 81 (2): 220-229. ]
- [48] Wang T G, Li M J, Wang C J, et al. Organic molecular evidence in the Late Neoproterozoic tillites for a palaeo-oceanic environment during the Snowball Earth era in the Yangtze region, southern China[J]. Precambrian Research, 2008, 162(3/4): 317-326.
- [49] Riedman L A, Porter S M, Halverson G P, et al. Organic-walled microfossil assemblages from glacial and interglacial Neoproterozoic units of Australia and Svalbard[J]. Geology, 2014, 42 (11): 1011-1014.
- [50] McKirdy D M, Burgess J M, Lemon N M, et al. A chemostratigraphic overview of the Late Cryogenian interglacial sequence in the Adelaide Fold-Thrust Belt, South Australia[J]. Precambrian Research, 2001, 106(1/2): 149-186.
- [51] Ma H R, Shen B, Lang X G, et al. Active biogeochemical cycles during the Marinoan global glaciation[J]. Geochimica et Cosmochimica Acta, 2022, 321: 155-169.
- [52] Hurtgen M T, Arthur M A, Suits N S, et al. The sulfur isotopic composition of Neoproterozoic seawater sulfate: Implications for a Snowball Earth? [J]. Earth and Planetary Science Letters, 2002, 203(1): 413-429.
- [53] 裴浩翔,李延河,付勇,等.贵州铜仁高地"大塘坡式"锰矿的成矿机制:硫,碳同位素制约[J]. 地球学报,2020,41(5):651-662. [Pei Haoxiang, Li Yanhe, Fu Yong, et al. Metallogenic mechanism of "Datangpo type" manganese deposits in Gaodi, Guizhou province: Constrains from sulfur and carbon isotopes [J]. Acta Geoscientica Sinica, 2020, 41(5): 651-662. ]
- [54] 满玲,祝圣贤,邓宾,等. 斯图特雪球地球存在持续的海洋碳 循环[J]. 沉积学报, 2024, 42 (1): 29-38. [Man Ling, Zhu Shengxian, Deng Bin, et al. Active marine carbon cycle during Sturtian "Snowball Earth" glaciation[J]. Acta Sedimentologica Sinica, 2024, 42(1): 29-38. ]
- [55] Hood A V S, Penman D E, Lechte M A, et al. Neoproterozoic syn-glacial carbonate precipitation and implications for a snowball Earth[J]. Geobiology, 2022, 20(2): 175-193.
- [56] Halverson G P, Maloof A C, Hoffman P F. The Marinoan glaciation (Neoproterozoic) in northeast Svalbard[J]. Basin Research, 2004, 16(3): 297-324.
- [57] Hoffman P F, Halverson G P, Schrag D P, et al. Snowballs in Af-

rica: Sectioning a long-lived Neoproterozoic carbonate platform and its bathyal foreslope (NW Namibia)[J]. Earth-Science Reviews, 2021, 219: 103616.

## Negative Carbon Isotope Excursions of the Statian Glacial Sediments in the Neoproterozoic Nanhua Basin and Their Geological Significance

WANG Ping<sup>1</sup>, WANG Jian<sup>1</sup>, DU YuanSheng<sup>2</sup>, ZHOU Qi<sup>3</sup>, YU WenChao<sup>2</sup>, XU Yuan<sup>2</sup>, YUAN LiangJun<sup>4</sup>, PAN Wen<sup>4</sup>, XU YaJun<sup>2</sup>

1. School of Resource and Environment, Henan Polytechnic University, Jiaozuo, Henan 454003, China

2. School of Earth Sciences, China University of Geosciences (Wuhan), Wuhan 430074, China

3. Engineering Technology Innovation Center of Mineral Resources Explorations in Bedrock Zones, Ministry of Natural Resources, Guiyang 550081, China

4. 103 Geological Party, Guizhou Bureau of Geology and Mineral Exploration and Development, Tongren, Guizhou 554300, China

Abstract: [Objective] The Neoproterozoic Struttin Glaciation (~717-660 Ma) developed widely across the world and was well recorded in the Nanhua Basin, but the study of carbon cycling during this period is lacking. To investigate the carbon cycle process during the Sturtian Glaciation, the carbon isotope compositions of the Tiesi'ao Formation precipitated during the glacial interval were systematically studied. [Methods] This study focuses on drillcore ZK2115, located in Gaodi Manganese Deposit, Songtao area, eastern Guizhou province, China. We analyzed highresolution organic and inorganic carbon isotopes  $(\delta^{13}C_{_{org}} \text{ and } \delta^{13}C_{_{carb}})$  from the syn-Sturtian Tiesi'ao Formation. **[Results]** The results show that the Tiesi'ao Formation is characterized by negative carbon isotope excursions, ranging between -9.29% and -3.37% (mean -7.24%). The  $\delta^{13}C_{org}$  values vary between -33.63% and -23.35% (mean -29.29‰). There is a positive correlation between the  $\delta^{13}C_{ev}$  and  $\delta^{13}C_{ev}$  values, indicating that the inorganic carbon isotope compositions are not affected by diagenesis and can be used to reflect the original carbon isotope signals of the Dissolved Inorganic Carbon (DIC) reservoir in the Nanhua Basin. Organic matter preserved in the Tiesi'ao Formation was generated through photosynthesis, and the carbon fixed in the organic matter was sourced from the DIC reservoir in seawater. [Conclusions] Combined with the low Total Organic Carbon (TOC) content (mean 0.2%) of the Tiesi'ao Formation, our results indicate that photosynthesis still occurred under the extreme glacial conditions at a low rate, although only a small amount of organic matter was generated. The study of carbon isotope compositions in glacial sediments can be used to explore carbon cycling during the Sturtian Glaciation, and to provide evidence for the paleomarine environment and biogeochemical cycle under extreme climatic conditions.

Key words: Sturtian Glaciation; Tiesi'ao Formation; carbon isotope; Nanhua Basin

<sup>[58]</sup> Swanson-Hysell N L, Rose C V, Calmet C C, et al. Cryogenian glaciation and the onset of carbon-isotope decoupling[J]. Science, 2010, 328(5978): 608-611.